

BB

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/28, 47/02, 47/12, 47/18	A1	(11) International Publication Number: WO 99/34821
		(43) International Publication Date: 15 July 1999 (15.07.99)
<p>(21) International Application Number: PCT/DK99/00006</p> <p>(22) International Filing Date: 6 January 1999 (06.01.99)</p> <p>(30) Priority Data: 98610001.4 9 January 1998 (09.01.98) EP</p> <p>(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).</p> <p>(72) Inventors: LANGBALLE, Peter; Efterårsvej 6, DK-2920 Charlottenlund (DK). NORUP, Elsebeth; Nordmarksvej 11, DK-4040 Jyllinge (DK).</p> <p>(74) Agent: PLOUGHMANN, VINGTOFT & PARTNERS A/S; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).</p>		<p>(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
(54) Title: STABILISED INSULIN COMPOSITIONS		
(57) Abstract		
<p>An aqueous insulin composition with improved chemical and physical stability comprising human insulin or an analogue or derivative thereof, a buffer selected from glycylglycine, citrate or TRIS, in particular glycylglycine, and metal ions, in particular calcium or magnesium ions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
ER	Estonia			SG	Singapore		

STABILISED INSULIN COMPOSITIONS

FIELD OF THE INVENTION

- 5 The present invention relates to stabilised pharmaceutical compositions comprising insulin or an analogue or derivative thereof. The invention also relates to parenteral formulations comprising such insulin compositions and to a method for improving the stability of insulin compositions.

10 BACKGROUND OF THE INVENTION

Diabetes mellitus is a metabolic disorder in which the ability to utilize glucose is more or less completely lost. About 2% of all people suffer from diabetes.

- 15 Since the discovery of insulin in the 1920's, continuous strides have been made to improve the treatment of diabetes mellitus. To help avoid extreme glucose levels, diabetic patients often practice insulin replacement therapy, whereby insulin is administered by injection.
- 20 In the treatment of diabetes mellitus, many varieties of insulin compositions have been suggested and used, including regular insulin, Semilente[®] insulin, isophane insulin, insulin zinc suspensions, protamine zinc insulin, and Ultralente[®] insulin. As diabetic patients typically are treated with insulin for several decades, there is a major need for safe and life quality improving insulin compositions. Some of the commercially available insulin
- 25 compositions are characterized by a fast onset of action, while other compositions have a relatively slow onset but show a more or less prolonged action. Fast acting insulin compositions are usually solutions of insulin, while retarded acting insulin compositions can be suspensions containing insulin in crystalline and/or amorphous form precipitated by addition of zinc salts alone or by addition of protamine or by a combination of both. In
- 30 addition, some patients use compositions having both a fast onset of action and a more prolonged action. Such a composition may be an insulin solution wherein protamine insulin crystals are suspended. Some patients prepare the final composition themselves by mixing an insulin solution with a suspension composition in the desired ratio.

Human insulin consists of two polypeptide chains, the so-called A and B chains, which contain 21 and 30 amino acid residues, respectively. The A and B chains are interconnected by two cystine disulphide bridges. Insulin from most other species has a similar construction, but may not contain the same amino acid residues at corresponding
5 positions.

The development of genetic engineering has made it possible to easily prepare a great variety of insulin compounds analogous to human insulin. In these insulin analogues, one or more of the amino acid residues have been substituted with other amino acid residues
10 which can be coded for by the nucleotide sequences. Since human insulin, as explained above, contains 51 amino acid residues, it is obvious that a large number of insulin analogues is possible, and a great variety of analogues with interesting properties have been prepared. In human insulin solutions with a concentration of interest for injectable compositions, the insulin molecule is present in associated form as a hexamer (Brange et
15 al. *Diabetes Care* 13, (1990), 923 - 954). After subcutaneous injection, it is believed that the rate of absorption by the blood stream is dependent on the size of the molecule, and it has been found that insulin analogues with amino acid residue substitutions which counteract or inhibit this hexamer formation have an unusually fast onset of action (Brange et al.: *Ibid*). This can be of great therapeutic value for the diabetic patient.

20

A general survey of pharmaceutical compositions containing insulin is given by Brange et al. in *Galenics of Insulin*, Springer-Verlag (Berlin, 1987).

Pharmaceutical compositions which are based on analogues of human insulin have e.g.
25 been presented by Heinemann et al., Luttermann et al. and Wiefels et al. at the "Frontiers in Insulin Pharmacology" International Symposium in Hamburg, 1992.

US 5 474 978 discloses a rapidly acting parenteral formulation comprising a human insulin analogue hexamer complex consisting of six monomeric insulin analogues, zinc
30 ions and at least three molecules of a phenolic derivative.

Normally, insulin compositions are administered by subcutaneous injection. What is important for the patient is the profile of action of the insulin composition, i.e. the action of insulin on the glucose metabolism as a function of the time from the injection,

including the time for the onset of insulin action, the maximum value and the total duration of action. A variety of insulin compositions with different profiles of action are required by patients. An individual patient may thus on the same day use insulin compositions with very different profiles of action. The profile of action required for any
5 given patient at any given time depends upon several factors, e.g. the time of the day and the amount and composition of any meal eaten by the patient.

Also important for the patient is the chemical stability of the insulin compositions, especially due to the abundant use of pen-like injection devices such as devices which
10 contain Penfill® cartridges, in which an insulin composition is stored until the entire cartridge is empty. This may last 1 to 2 weeks or more for devices containing a 1.5 or 3.0 ml cartridge. During storage, covalent chemical changes in the insulin structure occur. This may lead to the formation of molecules which are less active and potentially immunogenic such as deamidation products and higher molecular weight transformation
15 products (dimers, polymers, etc.). A comprehensive study on the chemical stability of insulin is given in by Jens Brange in "Stability of Insulin", Kluwer Academic Publishers, 1994.

Compositions comprising insulin and insulin analogues are traditionally formulated
20 using various additives, for example sodium phosphate (buffer), Zn^{2+} (stabiliser), phenol/m-cresol (preservative and stabiliser), sodium chloride (tonicity agent and stabiliser), and glycerol/mannitol (tonicity agents).

The shelf-life of insulin products is mainly compromised by the formation of soluble
25 aggregates (dimers and polymers) over time, despite the fact that insulin is typically stored at a low temperature of no more than about 5°C, which improves the shelf-life considerably compared to storage e.g. at room temperature. In addition, insulin products are subject to the formation of insoluble aggregates (fibrils) as a result of shaking, e.g. when carried in the pocket of a patient or during transport. It is essential
30 for the quality of an insulin product that the tendency to form such soluble and insoluble aggregates as a result of chemical or physical influences is reduced to an absolute minimum.

Although progress has been made in the chemical and physical stabilisation of insulin-containing compositions, the need still remains for improving the shelf-life of such products as well as their in-use time at temperatures of about 20-37°C, i.e. from room temperature to body temperature.

5

Acta Pharmaceutica Nordica 4(4), 1992, pp. 149-158 discloses insulin compositions with a sodium chloride concentration in the range of 0 to 250 mM. The major part of the compositions, including those which additionally comprise glycerol, contain a rather high amount of sodium chloride, i.e. 0.7%, corresponding approximately to a concentration of

10 120 mM.

PCT/DK97/00268 (Novo Nordisk A/S) discloses insulin compositions having improved chemical stability, the compositions comprising a combination of glycerol and/or mannitol and 5-100 mM of a halogenide, e.g. sodium chloride.

15

Storm & Dunn ("The Glu(B13) carboxylates of the insulin hexamer form a cage for Cd^{2+} and Ca^{2+} ions", *Biochemistry* 1985, 24, 1749-1756) performed studies showing that the highly co-operative zinc-insulin hexamer $(\text{In})_6(\text{Zn}^{2+})_2$ is further modulated by calcium binding, because the six Glu-(B13) residues within the hexamer structure form

20 a cage that binds calcium and cadmium ions. The single binding site is stated to bind Ca^{2+} and Cd^{2+} with high affinity and to be specific for these two ions.

SUMMARY OF THE INVENTION

- 25 It has now surprisingly been found that the stability of insulin compositions can be significantly improved by formulating the compositions using a combination of a buffer such as glycylglycine (Gly-Gly) and metal ions such as Ca^{2+} . It has in particular been found that when the sodium phosphate buffer in a traditional insulin composition is replaced with a Gly-Gly buffer in combination with calcium ions, the formation of
- 30 soluble aggregates during storage at 5°C decreased remarkably. This is especially surprising in view of the fact that a separate replacement of sodium phosphate by Gly-Gly was not found to have any effect, and that a separate addition of calcium to a composition containing a sodium phosphate buffer only had a minor effect. It was

further found that compositions containing the combination of Gly-Gly and calcium also showed a reduced tendency to form insoluble aggregates during shaking.

- It is an object of the present invention to provide pharmaceutical compositions
- 5 comprising insulin or an analogue or derivative thereof which have improved chemical and physical stability compared to known insulin compositions.

- In one aspect, the invention therefore relates to an aqueous insulin composition comprising human insulin or an analogue or derivative thereof, at least one buffer
- 10 selected from the group consisting of glycylglycine, citrate and TRIS, and metal ions selected from the group consisting of calcium and magnesium, with the proviso that when the buffer is TRIS, the concentration of metal ions is not in the range of 0.0004-0.01 M.

- 15 Another aspect of the invention relates to a parenteral pharmaceutical formulation comprising an insulin composition of the invention as defined above.

- A further aspect of the invention relates to a method for improving the stability of an insulin composition comprising human insulin or an analogue or a derivative thereof, the
- 20 method comprising adding to said composition at least one buffer selected from the group consisting of glycylglycine, citrate and TRIS, and metal ions selected from the group consisting of calcium and magnesium, with the proviso that when the buffer is TRIS, the concentration of metal ions is not in the range of 0.0004-0.01 M.

25 DETAILED DESCRIPTION OF THE INVENTION

Unless otherwise specified, the term "insulin" as used herein is intended to refer to not only human insulin as such, but also insulin analogues and derivatives.

- 30 The term "analogue of human insulin" as used herein refers to a polypeptide having the amino acid residue sequence of human insulin in which one or more amino acid residues have been deleted and/or replaced by other amino acid residues, including amino acid residues not encoded by the genetic code, or comprising additional amino acid residues, i.e. more than the 51 amino acid residues of human insulin.

The term "derivative of human insulin" as used herein refers to human insulin or an analogue thereof in which at least one organic substituent is bound to one or more of the amino acid residues.

5

The term "aqueous" as used herein refers to the fact that the insulin compositions of the present invention will be water-based, but it will be clear that the compositions may optionally contain additional solvents, e.g. a small amount of a water-miscible solvent.

10

The term "chemical stability" as used herein refers to the tendency of an insulin composition to form soluble aggregates of insulin during storage under static conditions, including storage at low temperatures of approximately 4-5°C. The term "physical stability" as used herein refers to the tendency of an insulin composition to

15 form insoluble aggregates of insulin as a result of a physical action such as shaking of an insulin composition.

In the present context, the unit "U" corresponds to 6 nmol of insulin.

20 Insulin compositions according to the invention have been shown to have a high chemical and physical stability, which is reflected in a reduction in the formation of soluble and insoluble aggregates after storage and/or shaking.

The present invention is particularly advantageous in connection with compositions
25 comprising analogues and/or derivatives of human insulin. Thus, the insulin composition according to the invention preferably comprises one or more fast-acting analogues of human insulin, in particular analogues wherein the amino acid residue at position B28 is Asp, Lys, Leu, Val or Ala and the amino acid residue at position B29 is Lys or Pro; or des(B28-B30), des(B27) or des(B30) human insulin. The insulin analogue is preferably
30 selected from analogues of human insulin wherein the amino acid residue at position B28 is Asp or Lys, and the amino acid residue at position B29 is Lys or Pro. The most preferred analogues are Asp^{B28} human insulin and Lys^{B28}Pro^{B29} human insulin.

In another embodiment the insulin composition according to the invention comprises an insulin derivative having a protracted profile of action, such an insulin having one or more lipophilic substituents. The preferred lipophilic insulins are acylated insulins, including those described in WO 95/07931 (Novo Nordisk A/S), e.g. human insulin derivatives
 5 wherein the ϵ -amino group of Lys^{B29} contains an acyl substituent which comprises at least 6 carbon atoms.

The following are preferred insulin derivatives:

N^{εB29}-myristoyl-des(B30) human insulin, N^{εB29}-palmitoyl-des(B30) human insulin, N^{εB29}-
 10 myristoyl human insulin, N^{εB29}-palmitoyl human insulin, N^{εB28}-myristoyl Lys^{B28} Pro^{B29} human insulin, N^{εB29}-palmitoyl Lys^{B28} Pro^{B29} human insulin, N^{εB30}-myristoyl-Thr^{B29} Lys^{B30} human insulin, N^{εB29}-palmitoyl-Thr^{B29} Lys^{B30} human insulin, N^{εB29}-(N-palmitoyl-γ-glutamyl)-des(B30) human insulin, N^{εB29}-(N-lithocholyl-γ-glutamyl)-des(B30) human insulin, N^{εB29}-(ω-carboxyheptadecanoyl)-des(B30) human insulin, and N^{εB29}-(ω-
 15 carboxyheptadecanoyl) human insulin; the most preferred being N^{εB29}-myristoyl-des(B30) human insulin.

Insulin compositions according to the invention will normally contain about 60 to 3000 nmol/ml, preferably 240 to 1200 nmol/ml, of human insulin or insulin analogue or
 20 derivative.

The amount of metal ions to be incorporated into compositions according to the invention is typically in the range of about 0.1-10, more preferably about 0.5-5, e.g. about 1-4 metal ions per hexamer of insulin or insulin analogue or derivative.
 25 Particularly good results have been obtained with about 2-3 metal ions per hexamer.

The metal ions should be pharmaceutically acceptable and are in particular calcium and/or magnesium. A preferred metal ion is calcium, as this has been shown to provide surprising and advantageous results in terms of stabilisation of insulin
 30 compositions containing Gly-Gly. However, it is also contemplated that other pharmaceutically acceptable metal ions having properties (size, valence) similar to calcium will also be suitable, in particular magnesium.

The concentration of the buffer, preferably Gly-Gly, will typically be in the range of about 1-20 mM, preferably about 3-15 mM, such as about 4-10 mM, e.g. about 5-7 mM.

- 5 In another embodiment, the insulin composition may comprise 5 to 100 mM, more preferably 10 to 100 mM, e.g. 10 to 70 mM, of a halide, as described in PCT/DK97/00268.

In a particular embodiment, the insulin composition of the invention comprises insulin or
10 an insulin analogue as well as an insulin derivative.

In a particular embodiment of the invention, the insulin composition comprises, in addition to metal ions and a buffer such as Gly-Gly as described above:

- 60 to 3000 nmol/ml, preferably 240 to 1200 nmol/ml, of human insulin or insulin
15 analogue or derivative,
10 to 40 µg Zn/100 U insulin, preferably 10 to 26 µg Zn/100 U insulin, and
0 to 5 mg/ml, preferably 0 to 4 mg/ml, of a phenolic compound.

- As a phenolic compound, 0.5 to 4.0 mg/ml, preferably 0.6 to 4.0 mg/ml, of m-cresol or
20 0.5 to 4.0 mg/ml, preferably 1.4 to 4.0 mg/ml, of phenol, or a mixture thereof, is advantageously employed.

- The insulin composition of the present invention may furthermore contain other ingredients common to insulin compositions, for example zinc complexing agents such as
25 citrate, or an additional buffer such as a phosphate buffer. When citrate is present, it may thus be present to function as a buffer and/or a zinc complexing agent.

The present invention furthermore relates to a parenteral pharmaceutical formulation comprising an insulin composition of the invention.

30

Moreover, the present invention is concerned with a method for improving the stability of an insulin composition comprising human insulin or an analogue or a derivative thereof, which method comprises adding at least one buffer selected from the group consisting of glycylglycine, citrate and TRIS and metal ions selected from the group consisting of

calcium and magnesium to said composition, with the proviso that when the buffer is TRIS, the concentration of metal ions is not in the range of 0.0004-0.01 M.

The invention is further illustrated by the following non-limiting examples.

5

EXAMPLE 1

Materials and methods

10

The composition of the test systems is given below; the buffer and calcium ion concentrations used are given in Table 1 below.

	Insulin ("NN304") *	0.6 μ mol
15	Phenol	1.50 mg
	m-Cresol	1.72 mg
	Mannitol	35 mg
	Sodium chloride	1.17 mg
	Zinc acetate	13.1 μ g Zn^{2+} (2 Zn^{2+} per (NN304) ₆)
20	Calcium chloride, dihydrate	see Table 1
	Buffer substance	see Table 1
	NaOH	for pH adjustment
	HCl	for pH adjustment
	Water to make	1 ml
25	pH = 7.5	

* NN304 is insulin analogue N⁶⁸²⁹-myristoyl-des(B30) human insulin.

Table 1. Concentration of buffer substance and Ca^{2+} for the test systems

Test system	di-Sodium hydrogen phosphate, dihydrate (mM)	Glycylglycin (mM)	Ca^{2+} ($\text{Ca}^{2+}/(\text{NN304})_0$)
Phosphate, 0 Ca^{2+}	7	0	0
Phosphate, 1 Ca^{2+}	7	0	1
Phosphate, 2 Ca^{2+}	7	0	2
Phosphate, 3 Ca^{2+}	7	0	3
Gly-Gly, 0 Ca^{2+}	0	7	0
Gly-Gly, 1 Ca^{2+}	0	7	1
Gly-Gly, 2 Ca^{2+}	0	7	2
Gly-Gly, 3 Ca^{2+}	0	7	3

5 Experimental procedure

50 ml of each test system were prepared. Insulin analogue NN304 was dissolved in water (2-10°C). Immediately after dissolution of NN304 solutions containing Zn^{2+} , Ca^{2+} , phenol + m-cresol + mannitol and buffer substance + NaCl were added. The pH value was adjusted by addition of HCl/NaOH, and water was added to make the final volume. The test systems were dispensed in 1.5 ml cartridges after sterile filtration (Sterivex GV 0.22 μ). The cartridges were placed at 4°C, 25°C or 37°C. Samples were taken and analysed for formation of dimers and polymers as well as for other impurities in the form of related proteins examined by liquid chromatography (referred to below as "RP impurities", i.e. impurities detectable by reverse phase HPLC). pH was also measured to determine the buffer effect. The analytical methods used in this study were Size Exclusion Chromatography and Reverse Phase Chromatography.

Results and discussion

Dimer formation

5 The values obtained for % dimer and the calculated values of k (% dimer/week) at 4, 25 and 37°C are listed in Table 2, 3 and 4, respectively. k is calculated by means of a linear regression. The listed values of R (correlation coefficient) indicate a rather good linear correlation between % dimer and storage time for the observation periods used. k is presented in the attached Figures 1, 2 and 3 as a function of the Ca^{2+} concentration.

The tables and figures show that the Ca^{2+} concentration has a greater effect on k (% dimer/week) for the Gly-Gly system compared to the phosphate system. The effect of Ca^{2+} is most pronounced at 4°C. An addition of 2-3 Ca^{2+} per $(\text{NN304})_6(\text{Zn}^{2+})_2$ thus causes a decrease of k to about one-third of the value without Ca^{2+} at 4°C for the Gly-Gly system, but only about a 20% decrease for the phosphate system. The rate of dimer formation is approximately identical for the two buffer systems without Ca^{2+} . The more pronounced effect of Ca^{2+} for the Gly-Gly system is believed to be due to a stronger binding between Ca^{2+} and phosphate than between Ca^{2+} and Gly-Gly. It is interesting to note that replacing phosphate with Gly-Gly causes a significant decrease of k at 25 and 37°C for a system without Ca^{2+} , but not at 4°C, which is the preferred storage temperature among the three temperatures tested.

Table 2. Formation of dimer at 4°C

Test system	Storage time (weeks)					k (% dimer/week)	R
	0 (% dimer)	10 (% dimer)	14 (% dimer)	18 (% dimer)	28 (% dimer)		
Phosphate 0 Ca ²⁺	0.19	0.29	0.30	0.36	0.46	0.00958	0.993
Phosphate 1 Ca ²⁺	0.17	0.26	0.27	0.33	0.42	0.00892	0.993
Phosphate 2 Ca ²⁺	0.18	0.26	0.26	0.33	0.40	0.00792	0.984
Phosphate 3 Ca ²⁺	0.17	0.23	0.26	0.31	0.38	0.00769	0.993
Gly-Gly 0 Ca ²⁺	0.17	0.30	0.30	0.38	0.46	0.0103	0.987
Gly-Gly 1 Ca ²⁺	0.19	0.23	0.26	0.30	0.36	0.00627	0.988
Gly-Gly 2 Ca ²⁺	0.18	0.21	0.21	0.24	0.27	0.00325	0.980
Gly-Gly 3 Ca ²⁺	0.17	0.19	0.20	0.24	0.27	0.00377	0.962

Table 3. Formation of dimer at 25°C

Test system	Storage time (weeks)				k (%dimer/week)	R
	0 (% dimer)	4 (% dimer)	6 (% dimer)	8 (% dimer)		
Phosphate 0 Ca ²⁺	0.19	0.36	0.49	0.58	0.0494	0.997
Phosphate 1 Ca ²⁺	0.17	0.34	0.41	0.52	0.0429	0.998
Phosphate 2 Ca ²⁺	0.18	0.34	0.46	0.52	0.0437	0.995
Phosphate 3 Ca ²⁺	0.17	0.34	0.46	0.50	0.0430	0.991
Gly-Gly 0 Ca ²⁺	0.17	0.26	0.39	0.43	0.0341	0.976
Gly-Gly 1 Ca ²⁺	0.19	0.24	0.34	0.40	0.0267	0.960
Gly-Gly 2 Ca ²⁺	0.18	0.22	0.29	0.36	0.0221	0.954
Gly-Gly 3 Ca ²⁺	0.17	0.21	0.29	0.35	0.0226	0.956

Table 4. Formation of dimer at 37°C

Test system	Storage time (weeks)				k (%dimer/week)	R
	0 (% dimer)	2 (% dimer)	4 (% dimer)	6 (% dimer)		
Phosphate 0 Ca ²⁺	0.19	0.54	0.81	1.12	0.153	0.999
Phosphate 1 Ca ²⁺	0.17	0.53	0.72	1.06	0.143	0.994
Phosphate 2 Ca ²⁺	0.18	0.49	0.74	1.06	0.145	0.999
Phosphate 3 Ca ²⁺	0.17	0.50	0.74	1.05	0.144	0.998
Gly-Gly 0 Ca ²⁺	0.17	0.33	0.56	0.88	0.118	0.989
Gly-Gly 1 Ca ²⁺	0.19	0.33	0.48	0.79	0.0975	0.979
Gly-Gly 2 Ca ²⁺	0.18	0.32	0.46	0.76	0.0940	0.979
Gly-Gly 3 Ca ²⁺	0.17	0.30	0.43	0.75	0.0935	0.970

5 Polymer

The values obtained for % polymer at 4, 25 and 37°C are listed in Tables 5, 6 and 7. % polymer as a function of time is presented in the attached Figure 4 at 37°C for the two buffer systems with 0 and 3 Ca²⁺/(NN304)₈(Zn²⁺)₂. Tables 5 and 6 show that the polymer formation is negligible at 4 and 25°C for the tested observation periods. The polymer formation is considerable at 37°C and is decreased by about 30% by addition of 2-3 Ca²⁺/(NN304)₈(Zn²⁺)₂ for both systems after 6 weeks of storage (Table 7 and Figure 4).

Table 5. Formation of polymer at 4°C

Test system	Storage time (weeks)				
	0	10	14	18	28
	(% polymer)	(% polymer)	(% polymer)	(% polymer)	(% polymer)
Phosphate 0 Ca ²⁺	0.09	0.06	0.07	0.07	0.09
Phosphate 1 Ca ²⁺	0.07	0.06	0.08	0.06	0.09
Phosphate 2 Ca ²⁺	0.07	0.06	0.08	0.06	0.07
Phosphate 3 Ca ²⁺	0.08	0.07	0.07	0.06	0.08
Gly-Gly 0 Ca ²⁺	0.06	0.05	0.07	0.08	0.12
Gly-Gly 1 Ca ²⁺	0.10	0.05	0.06	0.06	0.09
Gly-Gly 2 Ca ²⁺	0.08	0.06	0.07	0.06	0.08
Gly-Gly 3 Ca ²⁺	0.07	0.05	0.06	0.06	0.08

Table 6. Formation of polymer at 25°C

Test system	Storage time (weeks)			
	0 (% polymer)	4 (% polymer)	6 (% polymer)	8 (% polymer)
Phosphate 0 Ca ²⁺	0.09	0.09	0.15	0.09
Phosphate 1 Ca ²⁺	0.07	0.09	0.11	0.10
Phosphate 2 Ca ²⁺	0.07	0.09	0.11	0.09
Phosphate 3 Ca ²⁺	0.08	0.08	0.13	0.10
Gly-Gly 0 Ca ²⁺	0.06	0.10	0.11	0.13
Gly-Gly 1 Ca ²⁺	0.10	0.10	0.14	0.11
Gly-Gly 2 Ca ²⁺	0.08	0.10	0.11	0.10
Gly-Gly 3 Ca ²⁺	0.07	0.11	0.16	0.09

Table 7. Formation of polymer at 37°C

Test system	Storage time (weeks)			
	0 (% polymer)	2 (% polymer)	4 (% polymer)	6 (% polymer)
Phosphate 0 Ca ²⁺	0.09	0.32	0.30	0.53
Phosphate 1 Ca ²⁺	0.07	0.24	0.26	0.39
Phosphate 2 Ca ²⁺	0.07	0.25	0.16	0.35
Phosphate 3 Ca ²⁺	0.08	0.20	0.16	0.34
Gly-Gly 0 Ca ²⁺	0.06	0.24	0.29	0.53
Gly-Gly 1 Ca ²⁺	0.10	0.24	0.27	0.42
Gly-Gly 2 Ca ²⁺	0.08	0.17	0.21	0.38
Gly-Gly 3 Ca ²⁺	0.07	0.19	0.23	0.34

5 RP-purity

The values obtained for % impurities detectable by reverse phase HPLC at 4, 25 and 37°C are listed in Tables 8, 9 and 10. Ca²⁺ has no or perhaps only a small effect on the formation of RP-impurities. Table 10 shows a slightly increased amount of RP-10 impurities for the Gly-Gly systems compared to the phosphate systems after 6 weeks of storage at 37°C, but there are no apparent differences at 4 or 25°C.

Table 8. Formation of RP-impurities at 4°C

Test system	Storage time (weeks)	
	0 (% impurities)	10 (% impurities)
Phosphate 0 Ca ²⁺	2.4	2.3
Phosphate 1 Ca ²⁺	2.3	2.3
Phosphate 2 Ca ²⁺	2.4	2.3
Phosphate 3 Ca ²⁺	2.2	2.0
Gly-Gly 0 Ca ²⁺	2.3	2.3
Gly-Gly 1 Ca ²⁺	2.1	2.3
Gly-Gly 2 Ca ²⁺	2.3	2.1
Gly-Gly 3 Ca ²⁺	2.0	2.2

Table 9. Formation of RP-impurities at 25°C

Test system	Storage time (weeks)			
	0 (% impurities)	4 (% impurities)	6 (% impurities)	8 (% impurities)
Phosphate 0 Ca ²⁺	2.4	2.3	2.8	2.8
Phosphate 1 Ca ²⁺	2.3	2.4	2.5	2.9
Phosphate 2 Ca ²⁺	2.4	2.5	2.7	2.8
Phosphate 3 Ca ²⁺	2.2	2.5	2.9	2.8
Gly-Gly 0 Ca ²⁺	2.3	2.6	3.1	-
Gly-Gly 1 Ca ²⁺	2.1	2.5	2.9	3.0
Gly-Gly 2 Ca ²⁺	2.3	2.5	2.5	2.8
Gly-Gly 3 Ca ²⁺	2.0	2.4	2.6	2.5

Table 10. Formation of RP-impurities at 37°C

Test system	Storage time (weeks)			
	0 (% impurities)	2 (% impurities)	4 (% impurities)	6 (% impurities)
Phosphate 0 Ca ²⁺	2.4	3.0	2.9	3.7
Phosphate 1 Ca ²⁺	2.3	3.1	3.2	3.4
Phosphate 2 Ca ²⁺	2.4	3.0	3.2	3.6
Phosphate 3 Ca ²⁺	2.2	3.1	3.4	3.6
Gly-Gly 0 Ca ²⁺	2.3	3.4	3.7	4.5
Gly-Gly 1 Ca ²⁺	2.1	3.0	3.4	4.3
Gly-Gly 2 Ca ²⁺	2.3	2.9	3.1	4.0
Gly-Gly 3 Ca ²⁺	2.0	2.8	3.1	4.2

5

pH

The pH values of the phosphate and Gly-Gly buffers were compared at 25°C and 37°C, and it was found that the buffer effect of Gly-Gly was comparable to that of the phosphate buffer.

Conclusion

- The effect of Ca^{2+} on the dimer formation has been shown to be most pronounced for the Gly-Gly system at 4°C , compared to a phosphate buffer system. This is an important finding in light of the fact that insulin is typically stored at low temperatures of approximately $4-5^{\circ}\text{C}$. The addition of 2-3 Ca^{2+} per $(\text{NN304})_6(\text{Zn}^{2+})_2$ resulted in an approximately threefold decrease of the dimer formation for the Gly-Gly system, but only about a 20% decrease for the phosphate system at 4°C . The rate of dimer formation is approximately identical for the two buffer systems without Ca^{2+} at 4°C .
- The polymer formation was negligible at 4 and 25°C for the observation periods used. Polymer formation after 6 weeks at 37°C was decreased by about 30% by addition of 2-3 Ca^{2+} per $(\text{NN304})_6(\text{Zn}^{2+})_2$ for both the Gly-Gly and the phosphate system.
- Ca^{2+} has no or perhaps only a small effect on the formation of RP-impurities.

Based on the results presented above and the fact that formation of dimers and polymers constitutes a major chemical degradation pathway for insulin, it can be concluded that a combination of metal ions such as Ca^{2+} and a buffer such as Gly-Gly is advantageous as a stabiliser for aqueous insulin compositions.

EXAMPLE 2

- Tests were carried out in a similar manner as described above using "NN304" insulin (although a different batch than in Example 1), using either TRIS (7 mM) or Gly-Gly (7 mM) as the buffer and with the addition of either Mg^{2+} or Ca^{2+} as the metal ions. The test systems were prepared as described above and stored for a total of 48 weeks at a temperature of 5°C . The amount of dimer formation was determined after 0, 8, 16, 24, 32 and 48 weeks, and a k-value (% dimer/week) was determined using a linear regression. The systems tested and the results are shown in Table 11 below.

Table 11. Dimer formation in different systems at 5°C

Test system	Storage time (weeks)						k %dimer/week
	0 %dimer	8 %dimer	16 %dimer	24 %dimer	32 %dimer	48 %dimer	
Gly-Gly no metal ion	0.05	0.10	0.17	0.16	0.23	0.20	0.0033
TRIS no metal ion	0.05	0.09	0.10	0.14	0.17	0.16	0.0025
Gly-Gly 1 Mg ²⁺ / hexamer	0.05	0.08	0.09	0.10	0.14	0.12	0.0016
Gly-Gly 2 Mg ²⁺ / hexamer	0.04	0.06	0.09	0.08	0.11	0.10	0.0013
Gly-Gly 3 Mg ²⁺ / hexamer	0.04	0.06	0.09	0.10	0.13	0.10	0.0015
TRIS 2 Ca ²⁺ / hexamer	0.05	0.07	0.09	0.11	0.13	0.13	0.0018
TRIS 2 Mg ²⁺ / hexamer	0.05	0.08	0.08	0.10	0.13	0.12	0.0015

- 5 The results above show that the addition of Mg²⁺ to a Gly-Gly system results in a significant reduction of the rate of dimer formation. A reduction in the dimer formation is also seen by addition of Ca²⁺ or Mg²⁺ to the TRIS system, although the relative reduction in the TRIS system is smaller than for the Gly-Gly system. This example thus supports the conclusion above in Example 1, and further illustrates the advantageous
- 10 effect of using calcium or magnesium ions together with a buffer such as Gly-Gly for stabilising aqueous insulin compositions.

CLAIMS

1. An aqueous insulin composition comprising human insulin or an analogue or derivative thereof, at least one buffer selected from the group consisting of
5 glycylglycine, citrate and TRIS, and metal ions selected from the group consisting of calcium and magnesium,
with the proviso that when the buffer is TRIS, the concentration of metal ions is not in the range of 0.0004-0.01 M.
- 10 2. A composition according to claim 1 wherein the buffer is glycylglycine.
3. A composition according to claim 1 or 2 comprising 0.1-10, preferably 0.5-5, more preferably 1-4, e.g. 2-3 metal ions per hexamer of insulin or insulin analogue or derivative.
- 15 4. A composition according to any of claims 1-3, wherein the metal ions are calcium ions.
5. A composition according to any of claims 2-4 wherein the concentration of
20 glycylglycine is in the range of 1-20 mM, preferably 3-15 mM, such as 4-10 mM, e.g. 5-7 mM.
6. A composition according to any of the preceding claims, comprising an analogue of human insulin wherein the amino acid residue at position B28 is Asp, Lys, Leu, Val or Ala
25 and the amino acid residue at position B29 is Lys or Pro; or des(B28-B30), des(B27) or des(B30) human insulin.
7. A composition according to claim 6, comprising an analogue of human insulin wherein the amino acid residue at position B28 is Asp or Lys, and the amino acid residue at
30 position B29 is Lys or Pro, preferably Asp^{B28} human insulin or Lys^{B28}Pro^{B29} human insulin.
8. A composition according to claim 6, comprising des(B30) human insulin.

9. A composition according to any of claims 1-8, comprising a derivative of human insulin or an analogue thereof having one or more lipophilic substituents, preferably an acylated insulin, e.g. an insulin derivative wherein the ϵ -amino group of Lys^{B29} contains an acyl substituent comprising at least 6 carbon atoms.

5

10. A composition according to claim 9, wherein the insulin derivative is selected from the group consisting of N^{εB29}-myristoyl-des(B30) human insulin, N^{εB29}-palmitoyl-des(B30) human insulin, N^{εB29}-myristoyl human insulin, N^{εB29}-palmitoyl human insulin, N^{εB28}-myristoyl Lys^{B28} Pro^{B29} human insulin, N^{εB29}-palmitoyl Lys^{B28} Pro^{B29} human insulin, N^{εB30}-myristoyl-Thr^{B29}Lys^{B30} human insulin, N^{εB29}-palmitoyl-Thr^{B29}Lys^{B30} human insulin, N^{εB29}-(N-palmitoyl-γ-glutamyl)-des(B30) human insulin, N^{εB29}-(N-lithocholyl-γ-glutamyl)-des(B30) human insulin, N^{εB29}-(ω-carboxyheptadecanoyl)-des(B30) human insulin, and N^{εB29}-(ω-carboxyheptadecanoyl) human insulin.

15 11. A composition according to claim 10, wherein the insulin derivative is N^{εB29}-myristoyl-des(B30) human insulin.

12. A parenteral pharmaceutical formulation comprising an insulin composition according to any of the preceding claims.

20

13. A method for improving the stability of an insulin composition comprising human insulin or an analogue or a derivative thereof, which method comprises adding to said composition at least one buffer selected from the group consisting of glycylglycine, citrate and TRIS, and metal ions selected from the group consisting of calcium and

25 magnesium,

with the proviso that when the buffer is TRIS, the concentration of metal ions is not in the range of 0.0004-0.01 M..

14. A method according to claim 13, wherein the composition is formulated as defined in
30 any of claims 2-11.

1/2




Figure 1. k (%dimer/week) as a function of $\text{Ca}^{2+}/(\text{NN304})_6(\text{Zn}^{2+})_2$ at 4 °C.




Figure 2. k (%dimer/week) as a function of $\text{Ca}^{2+}/(\text{NN304})_6(\text{Zn}^{2+})_2$ at 25 °C.

2/2




Figure 3. k (%dimer/week) as a function of $\text{Ca}^{2+}/(\text{NN304})_6(\text{Zn}^{2+})_2$ at 37 °C.




Figure 4. %polymer as a function of time for the two buffer systems with 0 and 3 $\text{Ca}^{2+}/(\text{NN304})_6(\text{Zn}^{2+})_2$ at 37 °C.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/DK 99/00006

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/28 A61K47/02 A61K47/12 A61K47/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 472 385 A (BRANGE JENS J V ET AL) 18 September 1984 see the whole document	1,3, 12-14
A	WO 97 48414 A (NOVONORDISK AS) 24 December 1997 cited in the application	
A	WO 95 00550 A (NOVONORDISK AS ;BALSCHMIDT PER (DK)) 5 January 1995	
A	PATENT ABSTRACTS OF JAPAN vol. 014, no. 064 (P-1002), 6 February 1990 & JP 01 287463 A (KYOTO IKAGAKU KENKYUSHO:KK), 20 November 1989 see abstract	

	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 April 1999

Date of mailing of the international search report

21/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

FISCHER, W

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/DK 99/00006

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 681 833 A (DOTT RESEARCH LAB) 15 November 1995	

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/DK 99/00006

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4472385 A	18-09-1984	AT 23269 T	15-11-1986
		AU 550068 B	27-02-1986
		AU 8111482 A	16-09-1982
		CA 1170990 A	17-07-1984
		CH 652031 A	31-10-1985
		DK 100882 A,B,	11-09-1982
		EP 0060141 A	15-09-1982
		FI 820808 A,B,	11-09-1982
		FR 2501507 A	17-09-1982
		GB 2094145 A,B	15-09-1982
		IE 52904 B	13-04-1988
		BE 892413 A	08-09-1982
		JP 1707951 C	27-10-1992
		JP 3066290 B	16-10-1991
		JP 57209227 A	22-12-1982
		ZA 8201243 A	26-01-1983
WO 9748414 A	24-12-1997	AU 3253697 A	07-01-1998
		US 5866538 A	02-02-1999
WO 9500550 A	05-01-1995	AT 176482 T	15-02-1999
		AU 6995794 A	17-01-1995
		DE 69416409 D	18-03-1999
		EP 0705275 A	10-04-1996
		JP 2837956 B	16-12-1998
		JP 8511779 T	10-12-1996
		US 5547930 A	20-08-1996
		US 5834422 A	10-11-1998
		US 5840680 A	24-11-1998
EP 0681833 A	15-11-1995	JP 8027031 A	30-01-1996
		CN 1117874 A	06-03-1996
		US 5603943 A	18-02-1997